



復旦大學

Fudan University



復旦大學物理系物質科學報告

Physics Department Colloquium

New Opportunities for Spintronics: Magnetic Skyrmions

Dr. Axel Hoffmann

Materials Science Division, Argonne National Laboratory

The field of spintronics, or magnetic electronics, is maturing and giving rise to new subfields [1]. An important ingredient to the vitality of magnetism research in general is the large complexity due to competitions between interactions crossing many lengthscales and the interplay of magnetic degrees of freedom with charge (electric currents), phonon (heat), and photons (light) [2]. One perfect example, of the surprising new concepts being generated in magnetism research is the recent discovery of magnetic skyrmions. Magnetic skyrmions are topologically distinct spin textures that are stabilized by the interplay between applied magnetic fields, magnetic anisotropies, as well as symmetric and antisymmetric exchange interactions. Due to their topology magnetic skyrmions can be stable with quasi-particle like behavior, where they can be manipulated with very low electric currents. This makes them interesting for extreme low-power information technologies, where it is envisioned that data will be encoded in topological charges, instead of electronic charges as in conventional semiconducting devices. Towards the realization of this goal we demonstrated magnetic skyrmions in magnetic heterostructures stable at room temperature, which can be manipulated using spin Hall effects [3]. Furthermore, using inhomogeneous electric charge currents allows

the generation of skyrmions in a process that is remarkably similar to the droplet formation in surface-tension driven fluid flows [4]. However, detailed micromagnetic simulations show that depending on the electric current magnitude there are at least two regimes with different skyrmion formation mechanisms [5]. Lastly, we demonstrated that the topological charge gives rise to a transverse motion on the skyrmions, i.e., the skyrmion Hall effect, which is in analogy to the ordinary Hall effect originating from the motion of electrically charged particles in the presence of a magnetic field [6].

References

- [1] A. Hoffmann and S. D. Bader, Phys. Rev. Appl. **4**, 047001 (2015).
- [2] A. Hoffmann and H. Schultheiß, Curr. Opin. Solid State Mater. Sci. **19**, 253 (2015).
- [3] A. Hoffmann, IEEE Trans. Magn. **49**, 5172 (2013).
- [4] W. Jiang, *et al.*, Science **349**, 283 (2015).
- [5] O. Heinonen, *et al.*, Phys. Rev. B **93**, 094407 (2016).
- [6] W. Jiang, *et al.*, Nature Phys., doi:10.1038/nphys3883 (2016).

Time: 2:00pm, Tuesday, October 18, 2016

Location: Physics Building, Room 221B

(Cookies and coffee are served from 1:30 pm)